

PBM Laser Therapy Double-Blind Studies

Airaksinen O., et al.

Effects of infrared laser irradiation on the trigger points. *Skand J of Acu & EI Therapy*. 1988; 3: 56-61

Airaksinen O., et al.

Effects of laser irradiation on the treated and untreated trigger points. *Proc. 4th internal symposium. Acupuncture & Elektrother Res.* 1988; 13 (4): 238-239.

Antipa C. et al.

Clinical results of the low-energy laser action on post-traumatic nerve lesions in the distal forearm. *Laser therapy*. 1996; 1: 36.

Antipa C. et al.

Comparative effects of different IR low-energy diodes in the treatment of rheumatic diseases. 1997. In press (Monduzzi Editore, Bologna)

Armino L. et al.

Laser therapy for post-episiotomic neuralgia. *LASER. Journ Eur Med Laser Ass.* 1988; 1 (1): 7.

Atsumi K. et al.

Biostimulation effect of a low energy diode laser for pain relief. *Laser Surgery Med.* 1987; 7:77.

Barabas K. et al.

Controlled clinical and experimental studies on patients with rheumatoid arthritis and synovial membranes, carried out with neodymium phosphate glass laser radiation. *Proc. 7th Congress Internat Soc für Laserchirurgie und Med, Munich June 1987.* Abstract No. 216a.

Bihari I., Mester A., Abstract No. 216a.

The bio-stimulating effect of low-level laser therapy of a long-standing leg ulcer with helium-neon laser, helium-neon plus infrared laser, and incoherent light: preliminary report on a randomized, comparative double-blind study. *Laser therapy*. 1989; 1 (2): 97.

Boerner E. et al.

Double-blind study on the effectiveness of laser therapy. *SPIE Proc.* 1996. Volume 2929: 75-79.

Bihari I., Mester A.

The biostimulative effect of low-level laser therapy of a long-standing crural ulcer with helium-neon laser, helium-neon plus infrared laser, and incoherent light: preliminary report on a randomized, comparative double-blind study. *Laser therapy*. 1989; 1 (2): 97.

Carillo J. et al.

A double-blind, randomized, clinical study of the effectiveness of the helium-neon laser in

preventing pain, swelling and trismus after removal of an impacted third molar.
Int Dent Journ. 1990; 40:31.

Ceccherelli F.

Diode laser for cervical myofascial pain. A double-blind study compared to placebo.
The clinical J pain. 1989; 4: 301-304.

Cheng R.

Combined treatments of electrotherapy and soft laser therapy have a synergistic effect in relieving pain and curing the disease. Surgical and medical lasers.
1990; 3 (3): 135 (abstract).

Cieslar G. et al.

Effect of low power laser radiation in the treatment of musculoskeletal overload syndromes.
SPIE procedure. Volume 3198. 1997, pp. 76-82.

Cowen D. et al.

Low-energy helium-neon laser in the prevention of oral mucositis in patients undergoing a bone marrow transplant: results of a double-blind, randomized study.
Int J Radiat Oncol Biol Phys. 1997; 38 (4): 697-703.

de Bie RA

Effect of laser therapy on ankle sprains
Ned. T. Physiotherapy. 1988; 95: 108-112. (in Dutch)

Eckerdal A., Lehmann Bastian H.

Can low-reactive-level laser therapy be used in the treatment of neurogenic facial pain? A double-blind, placebo-controlled study of patients with trigeminal neuralgia.
Laser therapy. 1996; 8: 247-252.

Emmanoulidis O. et al.

The low power CW-IR laser application significantly accelerates the rehabilitation of competitive athletes with chronic pain. A double-blind study.
Laser Surgery Med. 1986; 6: 173.

England S. et al.

Low power laser therapy for shoulder tendinitis.
Skand J Rheumatology. 1989; 18: 427.

Flötter T., Refisch HP

Pain treatment with laser. A double-blind study. Proc. of the 4th boarding school symposium.
Acupuncture & Electro-Therapy Res. 1988; 13 (4): 236-237. Also: pain treatment with laser. A double-blind study. Cutting edge medicine. 1990; 4 (4): 52-56.

Fructuoso FJG, Moset JM

Estudio randomizado doble ciego sobre los efectos bioestimulantes del láser en la irradiación de glándula paratida en pacientes afectados de síndrome de Sjogren. (Double-blind study on the biostimulatory effects of laser irradiation on the parotid gland in patients with Sjogren's syndrome).
Examination and clinic laser. 1987; 4 (1): 18-25.

Gelskey SC et al.

The effectiveness of the Nd: YAG laser in treating dentin hypersensitivity.
J Can Dent Assoc. 1993; 59 (4): 337-386.

Gerschman JA et al.

Low level laser for dentin hypersensitivity.

Australian Dent J. 1994; 39: 6.

Goldman JA et al.

Laser therapy for rheumatoid arthritis.

Laser Surgery Med. 1980; 1: 93-102.

Gudmundsen J. et al.

Laser treatment for epicondylitis humeri and rotator cuff syndrome. Dobbelt blind study - 200 people. (Laser treatment of epicondylitis humeri and rotator cuff syndrome. Double-blind study - 200 patients. In Norwegian)

Norsk tidsskrift for idrettsmedisin. 1987; 2: 6.

Gertner C.

Low- Power Laser Analgesia (LPL): a controlled double-blind study in ankylosing spondarthritis (SPA).

Laser Surgery Med. 1989; Supplement 1:55.

Gärtte S. et al.

Double-blind study to test the efficacy and tolerability of low-energy laser therapy in patients with active osteoarthritis of the knee.

Jaros orthopedics. 1995. 12: 3034.

Haker E. et al.

Is low-energy laser treatment effective for lateral epicondylalgia? J pain and symptom management.

1991; 6 (4): 241.

Hashimoto K.

Clinical applications of various lasers in oral surgery.

Lasers in dentistry. Eds. Yamamoto Y et al. 1989; Pp. 63-70. Elsevier Science Publishing BV, Amsterdam

Hashimoto T. et al.

The effectiveness of laser irradiation on the area near the star ganglion is dose-dependent: a double-blind, crossover, placebo-controlled, crossover, double-blind study.

Laser therapy. 1997; 1 (9): 7-11.

Hopkins GO et al.

Double-blind crossover-crossover study of laser versus placebo in the treatment of tennis elbow. Proc International. Congress on lasers, "Laser Bologna".

1985: 210. Monduzzi Editore SpA, Bologna.

Hoteya K. et al.

Effects of a 1 W GaAlAs diode laser in the field of orthopedics. In: Conference Report: The First Congress of the International Society for Laser and Sports Medicine.

Tokyo, 1997. Laser Therapy 1997; 9 (4): 185.

Kaiser C. et al.

Estudio en doble ciego randomizado sobre la eficacia del He-Ne en el tratamiento de la sinusitis maxilar aguda: en pacientes con exacerbacion de una infeccion sinusal cronica. (Double-blind, randomized study of the effects of HeNe in the treatment of acute maxillary sinusitis: in patients

with exacerbation of chronic antrumitis).

Boleton CDL. 1986; 9:15.

Kamikawa K. et al.

Double-blind mid-laser experience in Japan. 1985. Proc Int Congress on Lasers, "Laser Bologna". 1985: 165-169. Monduzzi Editore SpA, Bologna.

Kemmotsu MD et al.

LLLT for pain relief - the current experience in the pain clinic. In: Advances in Laser Therapy. Eds Oshiro T, Calderhead R G. 1991: 197-200. John Wiley & Sons, Chichester, German ISBN 0-471-93154-3.

Khullar SM et al.

The low-level laser treatment improves long-lasting sensory aberrations of the inferior alveolar nerve after a surgical trauma.

J Oral Maxillofac Surg. 1996; 54: 2-7.

Khullar SM et al.

Effect of low-level laser treatment on neurosensory deficits after a sagittal Ramus split osteotomy. Oral surgery Oral medicine Oral pathology. 1966; 82 (2): 132-8.

Kim JW, Lee JO

Double-blind clinical cross-over study of an 830 nm diode laser and 5 years of clinical experience with biostimulation in plastic & aesthetic surgery in Asians.

Laser Surgery Med. 1998; Amendment 10:59.

Kinoshita F. et al.

Clinical evaluation of low-energy semiconductor laser therapy in oral surgery - a double-blind study. Josai Shika Daigaku Kiyo. 1986; 15 (3): 735-742. (in Japanese)

Laakso EL et al.

Pain scores and side effects in response to low-level laser therapy (LLLT) for myofascial trigger points.

Laser therapy. 1997; 2 (9): 67-72.

Lonauer G.

Controlled double-blind study on the effectiveness of He-Ne laser beams compared to He-Ne plus infrared laser beams in the therapy of activated arthrosis of the finger joints.

Clinical experiment rheumatism. 1987; 5 (suppl 2): 39

Longo L. et al.

Treatment with 904 nm and 10600 nm lasers of acute lumbago - double blind control.

LASER. Journ Eur Med Laser Ass. 1988; 1 (3): 16.

Lukas C. et al.

Low-level laser therapy for bedsores statium III.

Hoegschool van Amsterdam report. 1994.

Loegdberg-Andersson M. et al.

Low-level laser therapy (LLLT) for tendonitis and myofascial pain - a randomized, double-blind, controlled study.

Laser therapy. 1997; 2 (9): 79-86.

Do IT et al.

Helium-neon (red light) therapy for arthritis.

Rheumatology, 1983; 3:36.

Meier J.L., Kerkour K.
Traitement laser de la tendinite.
Med. And Hyg. 1989; 46: 907-911.

Mester A.
Biostimulating effect in wound healing by continuous wave laser diode 820 nm. Double-blind, randomized cross-over study.
Lasers in Med Science, Abstract Edition July 1988, No. 289.

Miyagi K.
Double-blind comparative study of the effect of low-energy laser radiation on rheumatoid arthritis.
Current awareness of Excerpta Medica. Amsterdam. Elsevier Science Publishers BV. 1989; 25: 315.

Mokhtar B. et al.
A double-blind, placebo-controlled study of the hypoalgesic effects of low-intensity laser irradiation of the roots of the neck using experimental ischemic pain. Proc.
Second meeting of the International Laser Therapy Assn., "London Laser", Sept. 1992, p. 61.

Mokhtar B. et al.
The Potential Impact of Pulse Repetition Rate in Laser-Mediated Analgesia: A Double-Blind, Placebo-Controlled Study Using Experimental Ischemic Pain. Proc.
Second meeting of the International Laser Therapy Assn, "London Laser" Sept 1992. p. 62

Molina JJ et al.
La laserterapia como coadyuvante en el tratamiento de la AR (Artritis Reumatoidea).
Boletin CDL, Barcelona. 1987; 14: 4-8.

Moore K. et al.
LLLT treatment for post-therapeutic neuralgia.
Laser therapy. 1988; 1: 7.

Moore K. et al.
The effect of infrared diode laser irradiation on the duration and severity of postoperative pain. A double-blind study.
Laser therapy. 1992; 4: 145.

Mousques T.
etude en double aveugle des effets du traitement unilateral au laser helium-neon lors de chirurgies parodontales bilaterales simultanées.
Search for Odonto-Stomatol. 1986; 11: 245.

Mousques T.
Etude en double aveugle des effets du helium-neon en chirurgie parodontale.
Find Odonto-Stomatol. 1986; 11: 223.

Neuman I. et al.
Low-energy phototherapy for allergic rhinitis and nasal polyposis.
Laser therapy. 1996. 1:37.

Nivbrant Bo et al.
Therapeutic laser treatment for osteoarthritis of the knee.
Acta-Orthop scandinavica. 1989; 60: 231.

Ortutay J et al.

Psoriatic arthritis treatment with low power laser radiation. A double-blind clinical study.
Laser medicine - Laser in Med Surg. 1998; 13 (3-4): 140.

Oyamada Y. et al.

A double-blind study of He-Ne laser therapy with low power lasers for rheumatoid arthritis.
Optoelectronics in Medicine. 1987; p 747-750. Springer Verlag, Berlin (excerpt). Full study in Boletón de CDL. 1988; 17: 8-12.

Palmgren N. et al.

Low power laser therapy for rheumatoid arthritis.
Lasers in medical science. 1989; 4: 193.

Palmgren N. et al.

Low-level laser therapy for infected abdominal wounds after surgery.
Laser Surgery Med. 1991; Amendment 3:11.

Palmieri B.

A double-blind, stratified, cross-over study of amateur tennis players with tennis elbows using infrared laser therapy.
Medical laser report. 1984; 1: 3-14

Rochkind S. et al.

Double-blind, randomized study of the use of neurotube and laser therapy in the treatment of complete sciatic nerve injury in rats.
Proc. 2nd Congr World Assoc. for Laser Therapy, Kansas City, 1998.

Roumeliotis D. et al.

820nm 15mW 4J / cm², laser diode application for sports injuries. A double-blind study.
Proc. British Medical Laser Association's fifth annual convention. 1987.

Saeki N. et al.

Double-blind test for biostimulation effects on pain relief from diode lasers.
1989. laser surgery; 1066: 93-100.

Sasaki K. et al.

A double-blind controlled study to analyze free amino acids in CO₂ laser burn wounds in a mouse model after doses of low-incident infrared (830 nm) diode laser energy.
Proc. 2nd meeting of the Internat Laser Therapy Assn., London, 1992, p.4.

Sasaki K. et al.

A preliminary double-blind controlled study to analyze free amino acids in burn wounds in mice after 830 nm diode laser therapy.
Laser therapy. 1997; 2 (9): 59-65.

Sato K. et al.

A double-blind evaluation of low power laser therapy in the treatment of post-therapeutic neuralgia. Surgical and medical lasers.
1990; 3 (3): 134 (summary)

Saunders L.

The effectiveness of low-level laser therapy for supraspinatus tendonitis.
Clinical rehab. 1995; 9: 126-134

Schindl A. et al.

Low-intensity laser radiation improves blood circulation in the skin in patients with diabetic microangiopathy.

Laser Surgery Med. 1998; Amendment 10: 7.

Scudds RA et al:

A Double-Blind Crossover Study of the Effects of a Low Power Gallium Arsenide Laser on Symptoms of Fibrositis.

Physiotherapy Canada. 1989; 41: (suppl. 3): 2.

Simunovic Z., Trobonjaca T. et al.

Treatment of medial and lateral epicondylitis - tennis and golf elbow - with low-level laser therapy: a multicenter, double-blind, placebo-controlled clinical study in 324 patients.

J Clin Laser Med & Surg. 1998; 16 (3): 145-151.

Simunovic Z., Trobonjaca T.

Soft tissue injuries during sports activities and traffic accidents - treatment with low-level laser therapy. A multicenter, double-blind, placebo-controlled clinical study in 132 patients.

Proc. IXX ASLMS Congress, Orlando, Florida, April 1999.

Snyder-Mackler L. et al.

Effect of the helium-neon laser on musculoskeletal trigger points.

Physical therapy. 1986; 66: 1087.

Snyder-Mackler L. et al.

Effect of helium-neon laser irradiation on peripheral sensory nerve latency.

Physical therapy. 1988; 68: 223.

Snyder-Mackler L. et al.

Effect of helium-neon laser radiation on skin resistance and pain in patients with trigger points in the neck or back.

Physical therapy. 1989; 69: 336.

Soriano FA et al.

Acute cervical spine pain is relieved with gallium arsenide (GaAs) laser radiation. A double-blind preliminary study.

Laser therapy. 1996; 8: 149-154.

Soriano FA et al.

Low response to laser therapy in patients with chronic lower back pain. A double-blind study.

Laser Surgery Med. 1998, supplement. 10, p. 6th

Toya S. et al.

Report on a computer-randomized double-blind clinical study to determine the effectiveness of the GaAlAs (830 nm) diode laser for pain relief in selected pain conditions.

Laser therapy 1994; 6: 143.

Taguchi T. et al.

Thermographic changes after laser irradiation in pain.

Clinical Laser Med Surg. 1991; 2 (9): 143.

Tsurko V. et al.

Laser therapy for rheumatoid arthritis. A clinical and morphological study.
Terap ark. 1983; 97. (Russian).

Volez-Gonzalez M. et al.

Treatment of relapse in herpes simplex in the lip and face area and primary herpes simplex in the genital area and the "area pudenda" with HeNe laser or acyclovir low power when administered orally.

SPIE Proc. 1995; Volume 2630: 43-50

Vasseljen O. et al.

Low-level lasers versus placebo in the treatment of tennis elbow. Scandal
Scandal J Reha Med. 1992; 24:37; also in physiotherapy. 1992; 5: 329.

Geher J.

Chronic pain relief with low power laser irradiation.

Neuroscientific Letters. 1983; 43: 339

Wanderer J.

Temporary suppression of clone in humans by brief photostimulation

. Brain research. 1985; 340: 109.

Walsh D. et al.

The effect of low-intensity laser radiation on stimulus conduction and skin temperature in the superficial radial nerve. Double-blind placebo-controlled study with experimental ischemic pain.
Proc. Second Boarding Meeting of the Laser Therapy Association, London, Sept. 1992.

Willner R. et al.

Low-power infrared laser biostimulation for chronic osteoarthritis of the hand.

Laser Surgery Med. 1985; 5: 149.

Wylie L. et al.

The hypoalgesic effects of low intensity infrared laser therapy on the mechanical pain threshold.

Laser Surgery Med. 1995; Amendment 7: 9.

Yamaguchi M. et al.

Clinical study of the treatment of hypersensitive dentin with a GaAlAs laser diode using the double-blind test.

Aichi Gakuin Daigaku Shigakkai Shi - Aichi-Gakuin Journal of Dental Science. 1990; 28 (2): 703-707. (in Japanese)

Abergel P. et al.

Control of connective tissue metabolism by laser: recent developments and future prospects.

Journal of the American Academy of Dermatology. 1984; 11: 1142.

Basko I.

A new frontier: laser therapy.

California Veterinarian. 1983; 10: 17.

Beck-Friis J., Borg G., Wetterberg L.

Rebound increase in nighttime melatonin levels after evening suppression by bright light exposure in healthy men: relationship to cortisol levels and morning exposure.

Wurtman RJ, ed. The Medical and Biological Effects of Light. Ann. NY Acad Sci. 1985; 453: 371-375.

Berki T. et al.

Biological effect of helium-neon (HeNe) laser irradiation with low power.
Lasers in medical science. 1988; 3:35.

Bihari I., Mester A.

The biostimulative effect of low-level laser therapy of a long-standing crural ulcer with helium-neon laser, helium-neon plus infrared laser and incoherent light: preliminary report of a randomized, comparative double-blind study.
Laser therapy. 1989; 1 (2): 97.

Bossy J. et al.

In-vitro investigation of the depth of penetration of low-energy laser beams in compact bones.
Faculte de Medecine and CHRU de Nimes, BP 26, 3000 NIMES, France. (1985).

Kalderkopf G.

Session Report

Ninth Congress of the International Society for Laser Surgery and Laser Medicine , Anaheim, California, USA: November 2-6 , 1991. Laser Therapy. 1992; 4 (1): 43.

Cherry R:

Measurement of protein rotation diffusion in membranes by flash photolysis.
Methods in Enzymology. 1978; (54): 47.

Derr VE et al:

Free radicals appear in some laser irradiated biological materials.
Federal proceedings. 1965; 24, No. 1, annex. 14: 99

Guang Hua Wang et al:

A study of the analgesic effect of low power HeNe lasers and their mechanism by electrophysiological means
Lasers in dentistry. Excerpta Medica. Elsevier Science Publishing House. 1989: p. 277.

Hachenberger I:

Laser beams for herpes diseases.
Medical cosmetology. 1981; 11: 142.

Haina D. et al:

Animal experiments on light-induced wound healing.
Proc from Laser-81, Opto-Electronics in Munich 1981.

Hong J. et al.: Animal experiments on light-induced wound healing:

Clinical study on laser therapy in 20 patients with post-therapeutic neuralgia.
Laser Therapy. 1990; 2 (4): 167.

Honmura A. et al:

Pain relieving effect of Ga-Al-As diode laser irradiation on hyperalgesia in carrageenin-induced inflammation.
Laser in Surgery Med. 1993; 13: 463.

Hort O, Vanpel T:

The distribution of Na⁺ and K⁺ under the influence of temperature gradients.
Pflügers Arch. 1971; 323: 158.

Horvath Z. et al:

Possible ab initio explanation of laser "biostimulation" effects.

Laser applications in medicine and surgery. Ed. G. Galetti et al: Proc 3rd World Congr - Intl Soc Low Power Laser Appln in Medicine 1992. Page 57.

KaihÅ,j P:

Low Level Lasers Effect pÅ ¥ FÅ,lsomme Tandhalse - en klinisk pilottest.

Odont practice. 1991; 6 (2): 229.

Karu T., Andreichuck T., Ryabykh T.

Suppression of chemiluminescence of human blood by diode laser irradiation at wavelengths of 660, 820, 880 or 950 nm.

Laser therapy. 1993; 5: 103.

Karu T:

Photobiological basics of low power laser therapy.

IEEE journal for quantum electronics. 1987; QE23 (10): 1703.

Karu T. and others:

Biostimulation of HeLa cells by low-intensity visible light.

Il Nuovo Cimento. 1982; Volume 1D, No. 6: 828.

Karu T. et al:

Biostimulation of HeLa cells by low-intensity visible light.

Il Nuovo Cimento. 1982; 1D (6): 828.

Kern's T:

HeNe lasers show promise in treating horse injuries.

Lasers and Applications. 1986; Dec: 39.

Kovacs I. et al:

Laser-induced stimulation of the vascularization of the healing wound.

Separatum-EXPERIENTIA. 1974; 30: 341

Kubota J, Ohshiro T:

The effects of low-reactive level laser therapy (LLLT) with diode laser on flap survival in a rat model.

Laser therapy. 1989; 1 (3): 127.

Kudoh Ch. Et al:

Effects of 830 nm gallium-aluminum-arsenide diode laser radiation on the saphenous nerve of the rat. Sodium-potassium adenosine-adenosine triphosphatase activity: a possible pain relief mechanism was investigated.

Laser therapy. 1989; 1 (2): 63. Landthaler M et al: Treatment of zoster, postherpetic pain and herpes simplex recurrences in loco with laser light. Progress Med. 1983; 101 (22): 1039.

Lubart R. et al:

A possible mechanism of the low-level laser-live cell interaction.

Laser therapy. 1990; 2 (2): 65.

Manne J:

Le laser arseniure de gallium 6 watts, etude clinique en odonto-stomatologie.

Le Chirurgien Dent de France 1985; 284: 15.

- Maricic B et al:
Analgesic effects of the laser in dental therapy.
Acta Stomat Croatian. 1987; 21 (4): 291.
- McKibbin L. and Paraschak D:
A study of the effects of lasers on chronically curved tendons at Whitney Hall Farm Limited, Canada,
January 1983.
Lasers in surgery and medicine. 1983; 3:55.
- Mester E. et al:
Studies on the inhibiting or promoting effect of the laser beams.
Arch Klin Chir. 1968; 322: 1022.
- Mester E. et al:
Effects of direct laser irradiation on human lymphocytes.
Arch Dermatol Res. 1978; 5:31
- Mester E. et al:
The biostimulating effect of the laser beam.
Proc from Laser - 81, Opto-Electronics in Munich 1981.
- Mizokami T. et al:
Effect of the diode laser on pain:
A clinical study of different types of pain. Laser therapy. 1990; 2 (4): 171.
- Montesinos M. et al:
Experimental effects of low-power lasers in enkephalin and endorphin synthesis.
LASER. Journ Eur Med Laser Ass. 1988; 1 (3): 2.
- Moore K. et al:
LLLT treatment of post-therapeutic neuralgia.
Laser Therapy. 1988; Pilot edition (1): 7.
- Muldiyarov P. et al:
Effect of monochromatic helium-neon laser red light on the morphology of zymosan arthritis in
rats. (Inst. For Rheumatism, Academy of Medical Sciences of the USSR, Mosc).
Biull Eksp Biol Med. 1983, Jan 95; 1: 55.
- Naeser M. et al.
Carpal tunnel syndrome - clinical result after low-level laser acupuncture, Microamps TENS and
other alternative therapies,
Journal of Alternative and Complementary Medicine Volume 4, No. 4, Nov. 1998
- Nasu F. et al:
Cytochemical effects of GaAlAs diode laser radiation on the activity of calcium ion-dependent
adenosine triphosphatase of the saphenous artery of the rat.
Laser therapy. 1989; 1 (2): 89.
- Oulamara A. et al:
Measurement of biological activity on botanical sample surfaces using a temporal decorrelation effect
of laser speckle.
Magazine for modern optics. 1989; 36 (2): 165.

Parascandolo S. et al:

Azione della Laser-terapia nella nevralgia essenziale del trigemino.

Int Congress on Lasers in Med and Surg, Bologna June 1985, p. 317. Monduzzi Editore SpA, Bologna, Italy.

Parrado C. et al:

Quantitative study of the morphological changes in the thyroid gland after IR laser radiation.

Lasers in Medical Sciences. 1990; 5: 77.

Passarella S. et al:

Increase of the electrochemical proton potential and the ATP synthesis in mitochondria of the rat liver, which were irradiated in vitro with a helium-neon laser.

FEBS letters. Sept 1984; 175 (1): 95.

Popowa M. et al:

Effect of the helium-neon laser beam on the regeneration of irradiated transplanted skeletal muscles.

Bulle Exp Biol Med. 1978; 80: 333. (ryska m eng abstr.)

Pourreau-Schnider N. et al:

The helium-neon laser treatment converts fibroblasts into myofibroblasts.

American Journal of Pathology. 1990; 137: 171.

Rochkind S. et al:

A single transcutaneous light irradiation of an injured peripheral nerve; comparative study of different wavelengths.

Read. in Med Sci. 1989; (4): 259.

Rochkind S. et al: A single transcutaneous light irradiation

Systemic effects of low-power laser irradiation on the peripheral and central nervous system, skin wounds and burns.

Lasers in surgery and medicine. 1989; 9: 174.

Rochkind S. et al:

Electrophysiological effect of the HeNe laser on the normal and injured sciatic nerve in the rat.

Acta Neurochir. (Vienna). 1986; 83: 125.

Shiroto C. et al:

Effects of diode laser radiation in vitro on the activity of human neutrophils.

Laser therapy. 1989; 1 (3): 135. Editore SpA, Bologna, Italy

Pääntinen P:

The effect of hair lasers on blood circulation in the skin.

Lasers in surgery and medicine Appendix 7, 1995, p. 9 (summary)

Shiroto C. et al:

Retrospective study of diode laser therapy for pain relief in 3635 patients:

detailed analysis using a questionnaire. Laser therapy. 1989; 1 (1): 41.

Wrench DC:

The active transport of water under temperature gradients.

Symp. Soc. Exp. Biol. 1954; 8:76.

Velez-Gonzalez M. et al:

Treatment of relapse in herpes simplex in the lip and face area and in primary herpes simplex in the genital area and "area pudenda" with a low-power HeNe laser or orally administered acyclovir

von Ahlften U et al:

SPIE Proc. 1995; Volume 2630-42: Experience in the treatment of aphthous and herpetiform diseases of the oral mucous membrane with a new infrared laser.

The bottom line. 1987; 5: 927.

Wakabayashi H. et al:

Effect of irradiation with a semiconductor laser on reactions that are induced in trigeminal caudal neurons by tooth pulp stimulation.

Laser in Surgery Med. 1993; 13: 605.

Walker J. et al:

Laser therapy for pain in trigeminal neuralgia.

Clinical J pain 1988; 3: 183.

Walker J:

Chronic Pain Relief from Low Power Laser Radiation.

Neuroscientific Letters. 1983; 43: 339.

Wakabayashi H. et al:

Treatment of dentin hypersensitivity using GaAlAs soft laser irradiation.

J Dent Res. 1988; 67: 182.

Wang L. et al:

A Review of Clinical Applications of Low Level Laser Therapy in Veterinary Medicine.

Laser therapy. 1989; 1 (4): 183.

Wilden L. and Dindinger D.:

Treatment of chronic complex diseases of the inner ear with low-level laser therapy in: Laser Therapy, Volume 8, 1996, No. 3, ISSN 0898-5901

Wilden L. and Karthein R.:

Import of radiation phenomena from electrons and therapeutic low-level lasers with regard to mitochondrial energy transfer in: Journal for Clinical Laser Medicine and Surgery, Volume 16, 1998, No. 3, ISSN 1044-5471

Wilden L. and Ellerbrock D.:

Improvement of hearing capacity through low-level laser light (LLLL) - observed using pre- and post-therapeutic audiometry courses of air and bone conduction over 12 frequencies of 0.125 Å ± 12 kHz in: Lasemedizin, Volume 14, 1999, No. 4, ISSN 0938-756X
